

HARMONIC ANALYSIS AND DYNAMICS FOR AFFINE ITERATED FUNCTION SYSTEMS

DORIN ERVIN DUTKAY AND PALLE E.T. JORGENSEN

ABSTRACT. We introduce a harmonic analysis for a class of affine iteration models in $\mathbb{R}^d$. Using Hilbert-space geometry, we develop a new duality notion for affine and contractive iterated function systems (IFSs) and we construct some identities for the Fourier transform of the measure corresponding to infinite Bernoulli convolutions.

CONTENTS

1. Introduction	1
2. Definitions	4
3. W -cycles for infinite Bernoulli convolutions	7
4. A harmonic function	9
5. λ representations	15
6. Strongly invariant measures	18
References	22

1. INTRODUCTION

There has been a recent increased interest in basis constructions outside the traditional setting of harmonic analysis. The setting which so far has proved more amenable to an explicit analysis with basis functions involves a mix of analysis and dynamics, and it typically goes beyond the standard and more familiar setting of orthonormal bases consisting of Fourier frequencies. The context of frames in Hilbert space (see, e.g., [ALTW04, BoPa05, BPS03]) is a case in point.

As is well known, the classical setting of Fourier presupposes an underlying group and Haar measure. But in dynamics, this is not available, and there is a variety of interesting cases which go beyond the group setting, and which nonetheless are amenable to harmonic basis constructions; see, e.g., [JoPe98, DuJo03, DuJo05]. These constructions are recursive in nature. At the same time, these recursive models arise in applications exhibiting a suitable *scale-similarity*. Their consideration combines classical ideas from infinite convolution with ideas from dynamics of a more recent vintage.

Work supported in part by the National Science Foundation.

2000 *Mathematics Subject Classification*. Primary 42A85; Secondary 42C25, 26A30, 28A80, 60G42, 47B80.

Key words and phrases. Measures, Fourier transform, projective limits, iterated function system (IFS), transfer operator, Hilbert space, Perron-Frobenius, harmonic function, cycle, Markov process, Bernoulli convolution.

In the simplest and still useful non-group case, the similarity may be defined by a scaling of vectors in $\mathbb{R}^d$ with a fixed d by d matrix. Our sets and measures then arise as attractors of an associated dynamical system. The sets and measures in our present non-group context arise from a certain algorithmic construction on a fixed finite system of affine mappings in $\mathbb{R}^d$. Consideration of bases in this context involves a new viewpoint and a combination of ideas from dynamics and harmonic analysis.

Our aim in this paper is to develop a notion of cyclicity from dynamical systems and operator theory, and to apply it to geometric iteration systems. Specifically, we create a harmonic analysis on the class of affine iterated function systems (IFS). Our present harmonic analysis is based on Hilbert-space geometry and a certain duality notion, but it is more versatile than the earlier such notions from fractal Fourier analysis; see especially [JoPe98]. As is well known, IFSs provide us with prototypes of fractals, and they have for some time served as successful models for symbolic dynamics, e.g., [BKS91]. More generally, fractals of the affine IFS-type arise in several areas in symbolic dynamics; see, e.g., [Edg98].

Specifically, we develop in this paper a concept of cyclicity which generalizes a harmonic analysis which was studied in earlier papers by the second named author, S. Pedersen, and R. Strichartz, among others. While it was shown in [JoPe98] that certain affine and contractive iterated function systems in $\mathbb{R}^d$ admit Hilbert-space-orthonormal bases (ONBs), it turns out that the conditions for existence of ONBs in affine IFSs is somewhat restrictive. The affine systems under discussion include Cantor sets of Sierpinski type, realized in $\mathbb{R}^d$.

In Section 2 below, we recall how a fixed iterated function system (IFS) $(\tau_i)_i$ gives rise to a unique attractor X . As a result, we get induced maps in X , also denoted by τ_i . We further recall how the index labels for the system (τ_i) introduce an encoding space Ω for the points in X . Using the natural shift map of Ω and the notion of periodicity, we may then introduce associated cycles C as subsets of X . A certain weight function W is fixed and defined on X ; and it allows us to introduce transition probabilities associated with (τ_i) , a transition operator R_W , and a path-space measure $P_x = P_x^{(W)}$. We further use W to define certain W -cycles which are key to understanding the solutions to the equation $R_W h = h$. These solutions h are called harmonic functions, or R_W -harmonic functions.

Our main results Theorems 4.6, 4.10 are about the harmonic functions for the special IFSs of Bernoulli type. Section 5 deals with some geometric issues used in Section 4. Section 6 is about general IFSs.

It turns out that the introduction of a harmonic analysis for these systems adds to their use, and it shows that seemingly similar Cantor sets can have drastically different harmonic analysis: For example, the usual Cantor subset of the line, arising from subdivision by 3, is vastly different from the corresponding Cantor construction based instead on subdivision by 4; see [JoPe98] for details. In this paper, we identify a notion of extremality for certain finite cycles in IFSs. Among these cycles, we identify a small subclass of cycles where the given weight function W attains its maximum. The choice of weight function W depends on and is intimately tied to the geometry of the IFS under discussion. In fact, W serves to create a duality of affine IFSs via a certain complex Hadamard matrix; see, e.g., [JoPe96]. When the IFS and W are specified, we call the corresponding extreme cycles W -cycles (Definition 2.3). We point out with examples the significance of these W -cycles;

and we find all the W -cycles for particular affine fractal systems. It turns out that the case when an affine IFS has the ONB property (such as the Cantor set arising by quarter partitions and skipping every second subinterval) corresponds to the situation in our present setup when the only extreme W -cycle is a one-cycle. This last fact holds more generally; but as we show, not all systems for which the only W -cycle is a one-cycle admit an ONB of Fourier type.

A W -cycle is a cycle C for (X, τ_i) such that $W(y) = 1$ for all y in C (Definition 2.3). The function W determines a transition operator R_W (Definition 2.2) and a random walk on a space of paths beginning at points in x . Hence, each x induces a Kolmogorov measure P_x on Ω (equation (2.7)). We show that each P_x is countably atomic, i.e., is supported on a countable set S in Ω , and each point in S is an atom for P_x (Corollary 4.7, Theorem 4.6). In particular, we give the precise structure of this P_x for a classical one-parameter family of infinite Bernoulli convolution measures ν_λ (Examples 3.1, 3.2), and associated support sets X_λ . Using this, we further show that several properties of ν_λ are determined directly from the pair (P_x, R_W) . For example, we find the support X_λ of ν_λ (Proposition 5.1); and we show that for all λ , the absolute square of the Fourier transform of ν_λ is the unique solution to a functional equation defined directly from P_x (Theorem 4.10). The discrete set S which supports the measures P_x encodes the union of the W -cycles (Lemma 4.1); and our proofs are based on a detailed analysis of the W -cycles for the duality of the Bernoulli systems. In particular, we find all the W -cycles for the Bernoulli systems and we show (Theorem 4.6) that there is a one-to-one correspondence between the W -cycles C and certain extreme solutions $h(x) = h_C(x) = P_x(\{C\text{-cyclic paths}\})$ to the eigenvalue problem $R_W h = h$, called harmonic functions (see Definition 2.5). We show that there is a specific set D of rational values of λ , such that if λ is in $(0, 1) \setminus D$, then $H_\lambda := \{h \in C(X_\lambda) \mid R_W h = h\}$ is one-dimensional, and if λ is in D , then $\dim H_\lambda = 2$. For all λ , we show that $\sum_C h_C = 1$ on X_λ .

When W is assumed normalized, we obtained in [DuJo05] that, under certain conditions, $\sum_C h_C = 1$, i.e., the constant function “one”. But we also gave examples in $\mathbb{R}^d$ when this does not hold. As an application of our W -cycles, we show that for the Bernoulli IFSs, this sum is the constant function one for *all* λ in $(0, 1)$. Here we address this issue and related questions for the harmonic analysis of Bernoulli IFSs, and our discussion is divided up into the two cases: (i) λ in $(0, \frac{1}{2})$, the fractal case, and (ii) $\lambda \geq \frac{1}{2}$. It is the second case that holds more surprises.

Our first result (see Theorem 3.3) for these Bernoulli systems is that when λ is in $[1/2, 1)$ there are no W -cycles of minimal period $p > 1$. If $\lambda > 1/2$, and $\lambda \notin D := \{1 - \frac{1}{2n} \mid n \in \mathbb{N}\}$, then the only W -cycle is $C = \{0\}$ as a subset of the interval X_λ which supports the system. The corresponding eigenfunction, or harmonic function, is then $h_0(x) = P_x(\mathbf{N}_0)$, and $\mathbf{N}_0$ is the set of paths that end in an infinite repetition of the trivial cycle $\{0\}$. So either h_0 is the constant function 1, or $\dim H_\lambda > 1$.

In Theorem 4.6, we show that if $\lambda > \frac{1}{2}$ falls in the complement of the countable and exceptional set D , then we may conclude that h_0 has no zeroes. Then by a theorem from [DuJo05], h_0 must be the constant function one, and indeed we get $\dim H_\lambda = 1$. The significance of this may be understood from our result, Theorem 4.10. This theorem shows that $|\hat{\nu}_\lambda|^2$ is the unique solution to a certain functional

equation, i.e., the modulus-square of the Fourier transform of the Bernoulli measure ν_λ is determined this way for all λ in the complement of D .

For λ in the exceptional set D , the space H_λ is also understood (see part two in Theorem 4.6): In that case, h_0 is not a constant function; the space H_λ is then 2-dimensional, and it is generated by two distinct W_λ -cycles C_0 and C_1 . Each of the two cycles defines a harmonic function, h_0 and h_1 , respectively. The two functions form a basis for H_λ , and $h_0 + h_1 = 1$.

2. DEFINITIONS

Definition 2.1. An affine IFS in $\mathbb{R}^d$ is determined by

- R : a $d \times d$ matrix such that the eigenvalues λ satisfy $|\lambda| > 1$;
- $B \subset \mathbb{R}^d$ a finite subset.

Then we define

$$(2.1) \quad \tau_b(x) := R^{-1}(x + b), \quad (x \in \mathbb{R}^d),$$

and note that there is a unique compact subset $X_B \subset \mathbb{R}^d$ such that

$$(2.2) \quad X_B = \bigcup_{b \in B} \tau_b(X_B).$$

In fact X_B consists of the points

$$(2.3) \quad x = \sum_{j=1}^{\infty} R^{-j} b_j, \quad \text{as } b = (b_1, b_2, \dots) \in \prod_{\mathbb{N}} B \quad (= B^{\mathbb{N}}).$$

Set $\Omega = \Omega_B = B^{\mathbb{N}}$;

- $\mathcal{W}$:= all finite words in B ;
- $\mathcal{W}_k := \{(b_1 \dots b_k) \mid b_i \in B\}$;
- If $b \in \mathcal{W}$, let $|b|$ denote the length of b , i.e., $|(b_1 \dots b_k)| = k$.

For $b \in \Omega_B$, set

$$(2.4) \quad \pi_R(b) := \sum_{j=1}^{\infty} R^{-j} b_j.$$

For $k \in \mathbb{N}$, and $b \in \mathcal{W}_k$, set

$$(2.5) \quad \pi_R(b) := \sum_{j=1}^k R^{-j} b_j.$$

When an affine IFS (X, τ_i) is given, our duality is then determined by a certain function $W: X \rightarrow [0, \infty)$ satisfying

$$(2.6) \quad \sum_{i=1}^N W(\tau_i x) = 1,$$

and by an associated transfer operator R_W acting on functions on X (Definition 2.2). We show that R_W has properties analogous to those of finite Perron-Frobenius non-negative matrices.

For a non-negative function W on X satisfying (2.6) following Kolmogorov, one can define probability measures P_x on Ω , $x \in X$, such that, for a function f on Ω which depends only on the first n coordinates,

$$(2.7) \quad \int_{\Omega} f(\omega) dP_x(\omega) = P_x(f) \\ = \sum_{\omega_1, \dots, \omega_n} W(\tau_{\omega_1} x) W(\tau_{\omega_2} \tau_{\omega_1} x) \cdots W(\tau_{\omega_n} \cdots \tau_{\omega_1} x) f(\omega_1, \dots, \omega_n).$$

We will assume throughout that the *normalization* (2.6) holds. In particular W_λ in (3.4) for the λ -Bernoulli system satisfies (2.6).

The main question addressed in this paper arose in, or is motivated by, a number of earlier investigations on iterated function systems (IFS); see, e.g., [Edg98], [JoPe96], [JoPe98], [So95], [PeSo96], [KSS03], and [Jor04]. For additional details on the measures P_x occurring in (2.7), see, for example, [Jor06]. The Bernoulli convolutions (see details below) are of interest in their own right, and serve as models for general structures in dynamics. They were studied in the thirties by P. Erdős [Er39] and subsequently by many others.

Infinite Bernoulli convolutions (Example 3.1, Proposition 5.1) form a subclass of contractive iterated function systems (IFS) of affine type (Definition 2.1). Each such affine IFS $(X, (\tau_i)_{i=1, \dots, N})$ has a unique measure ν_p on X which assigns prescribed probabilities p_i to the branch maps τ_i . We establish a duality for these measures, and we show in particular that each ν_p is non-atomic (Corollary 6.6).

Let λ be in the open interval $(0, 1)$. Let $N = 2$, $p_i = \frac{1}{2}$, let the two τ -maps be $\tau_{\pm}(x) := \lambda x \pm 1$, and denote the corresponding measure by ν_λ (Example 3.2). This measure ν_λ is the distribution of the random variable $x(\omega) = \sum_{k=1}^{\infty} \pm \lambda^k$ with independent and random assignment of $\pm$ signs. Moreover, ν_λ is an infinite convolution measure, studied first by P. Erdős, and is called a Bernoulli convolution. Let Ω be the space of all configurations ω of signs in x . We show that each Bernoulli system is *one side* in a duality of affine IFSs, and that this duality holds more generally for contractive affine IFSs.

The IFS $\tau_{\pm}(x) = \lambda x \pm 1$ defined for fixed λ is called a *Bernoulli IFS*; and for each λ in $(0, 1)$, there is a natural choice for the function $W = W_\lambda$, see equation (3.4), and an associated process P_x .

Definition 2.2. A third essential ingredient for the process is a transfer operator R_W depending on W . For functions f on X we set

$$(R_W f)(x) = \sum_{i=1}^N W(\tau_i x) f(\tau_i x) \quad (x \in X).$$

We denote the operator R_W , the *Ruelle operator*, or a transfer operator, to stress the dependency on the function W . Each R_W defines special *harmonic functions*

$$H = \{h \in C(X) \mid R_W h = h\}.$$

The main results in this paper concern the real part of the peripheral spectrum of this operator R_W ; see especially Theorem 4.6 below. But to state and prove our spectral results for R_W , we must first introduce a certain encoding map for our IFSs, and an associated family of finite cycles, called *W-cycles*.

Definition 2.3. [Cycles and W -cycles] If a point $x \in X$ has the property that $\tau_{\omega_p} \cdots \tau_{\omega_1} x = x$ for some $\omega_1, \dots, \omega_p \in \{1, \dots, N\}$, then the set

$$C := \{x, \tau_{\omega_1} x, \dots, \tau_{\omega_{p-1}} \cdots \tau_{\omega_1} x\}$$

is called a *cycle*. If p is minimal with this property, then p is called the length of the cycle and C is called a p -cycle. We can identify the cycle C with the finite word $\omega_1 \dots \omega_p$.

Every cycle with period p is also periodic with a period that is a multiple of p . We say that a cycle C has minimal period p if C does not close up with a period smaller than p , i.e., with a period that is a divisor in p . Unless specified otherwise, we will assume that the term p -cycle refers to the minimal period.

When W is given, we are interested in those cycles C satisfying $W(x) = 1$ for all x in C . These cycles are called *W -cycles*. And we will use the term *W - p -cycle* for a p -cycle which is also a W -cycle.

Definition 2.4. Let $\Omega = \{1, \dots, N\}^{\mathbb{N}}$. We say that Ω is an encoding of $(\tau_i)_{i=1}^N$ if for all $\omega = (\omega_1 \omega_2 \dots) \in \Omega$

$$(2.8) \quad \bigcap_{n=1}^{\infty} \tau_{\omega_1} \tau_{\omega_2} \cdots \tau_{\omega_n}(X)$$

is a singleton $\pi(\omega)$. (This condition is automatic if the maps τ_i are strictly contractive, but it is known to hold also for important classes of IFS which are not contractive.)

Suppose an IFS has an encoding π . Then $\pi: \Omega \rightarrow X$ maps onto the attractor of the IFS X_τ , and π is called the *encoding mapping*.

If $(\omega_1 \dots \omega_p)$ is a word in the alphabet $\{1, \dots, N\}$, then

$$(\omega_1 \dots \omega_p)^\infty = (\omega_1 \dots \omega_p \omega_1 \dots \omega_p \dots)$$

is the point in Ω resulting from an infinite repetition of $(\omega_1 \dots \omega_p)$. It generates a p -cycle under the shift σ in Ω , i.e.,

$$(2.9) \quad \sigma(\eta_1 \eta_2 \eta_3 \dots) := (\eta_2 \eta_3 \eta_4 \dots),$$

i.e.,

$$\sigma^p((\omega_1 \dots \omega_p)^\infty) = (\omega_1 \dots \omega_p)^\infty.$$

This p -cycle will be denoted $C(\omega_1 \dots \omega_p)$, and it is said to be generated by the word $(\omega_1 \dots \omega_p)$.

Definition 2.5. For each W -cycle C , consider the subset $\mathbf{N}_C$ in Ω consisting of paths that end up in C after a finite number of backwards shifts, i.e., $\mathbf{N}_C$ consists of all finite words, followed by an infinite repetition of the fixed word representing C . That is, if C is generated by the word $(k_1 \dots k_p)$, then

$$\mathbf{N}_C := \{\omega_1 \dots \omega_n (k_1 \dots k_p)^\infty \mid \omega_1, \dots, \omega_n \in \{1, \dots, N\}, n \in \mathbb{N}\}.$$

We define the function

$$(2.10) \quad h_C(x) = P_x(\mathbf{N}_C).$$

Then h_C is harmonic and continuous (see [DuJo05, Proposition 5.8, Remark 5.9]), and the question is if the space H of continuous harmonic functions is generated by the W -cycles.

Note the following distinct parts of the conclusion: First that the function $h_C(x) = P_x(\mathbf{N}_C)$ is continuous, and secondly, that it satisfies the equation $R_W h_C = h_C$. The reader is also referred to [DuJo05] for the complete arguments of this part. It turns out that properties for the functions h_C , such as existence of zeroes, can be checked; and that these properties in turn have implications both for the measures P_x and for the strongly invariant IFS measures ν .

3. W -CYCLES FOR INFINITE BERNOULLI CONVOLUTIONS

Example 3.1. Bernoulli's infinite convolution; see [SSU01], [So95], [PeSo96], [PeSo00], and [Er39]. Let $d = 1$. Let $\lambda \in (0, 1)$, and set $R = \lambda^{-1}$, $B = \{0, 1\}$. Then Ω_B is the usual probability space $\Omega_B = \{0, 1\}^{\mathbb{N}}$.

Example 3.2. A dual pair of Bernoulli systems.

Let $\lambda \in (0, 1)$ and set $R = \lambda^{-1}$. We consider IFSs corresponding to two sets

$$(3.1) \quad B(\lambda) := \{\pm\lambda^{-1}\}, \quad \text{and} \quad L(\lambda) := \left\{0, \frac{1}{4}\right\},$$

i.e.,

$$(3.2) \quad \tau_{\pm}(x) := \lambda x \pm 1;$$

and

$$(3.3) \quad \tau_0(x) := \lambda x, \quad \tau_{1/4}(x) := \lambda \left(x + \frac{1}{4}\right).$$

The choice of B , and the term *duality*, are motivated by the general theory of affine IFSs as developed in [JoPe96], [JoPe98], and [DuJo05], as well as in other recent papers on the harmonic analysis of IFSs.

Setting

$$(3.4) \quad W(x) := W_B(x) = \cos^2(2\pi\lambda^{-1}x),$$

and

$$(3.5) \quad (R_W f)(x) := W(\tau_0 x)f(\tau_0 x) + W(\tau_{1/4} x)f(\tau_{1/4} x),$$

we see that

$$(3.6) \quad R_W f(x) = \cos^2(2\pi x)f(\lambda x) + \sin^2(2\pi x)f(\lambda(x + 1/4)),$$

and

$$(3.7) \quad R_W \mathbf{1} = \mathbf{1},$$

where $\mathbf{1}$ denotes the constant function “one”.

We will see in Proposition 5.1 that if $\lambda \geq 1/2$, then the two attractors $X_{B(\lambda)}$ and $X_{L(\lambda)}$ consist of the following two intervals:

$$X_{B(\lambda)} = \left[\frac{-1}{1-\lambda}, \frac{1}{1-\lambda} \right] \quad \text{and} \quad X_{L(\lambda)} = \left[0, \frac{\lambda}{4(1-\lambda)} \right].$$

The classical theory of Bernoulli convolutions [PSS00] has a point of contact with the scaling equation [Dau92] from wavelet theory. To illustrate this point, consider functions f on $\mathbb{R}$, and the equation

$$(3.8) \quad f(x) = \frac{1}{2\lambda} \left(f\left(\frac{x-1}{\lambda}\right) + f\left(\frac{x+1}{\lambda}\right) \right).$$

Its integrated form is as follows:

$$(3.9) \quad F(x) = \frac{1}{2} \left(F\left(\frac{x-1}{\lambda}\right) + F\left(\frac{x+1}{\lambda}\right) \right).$$

Then, by elementary multi-scale theory, the special case of $\lambda = \frac{1}{2}$ in (3.8) coincides with Haar's scaling identity; see, e.g., [Dau92]. To see this, recall that Haar's scaling function is $f = \frac{1}{4}\chi_{[-2,2]}$. A calculation shows that (3.9) is solved by

$$F(x) = \begin{cases} 0 & \text{if } x < -2, \\ \frac{1}{4}(x+2) & \text{if } -2 \leq x < 2, \\ 1 & \text{if } 2 \leq x. \end{cases}$$

In general, if λ is in $(0, 1)$ and ν_λ denotes the Bernoulli measure, then the cumulative distribution function $F_\lambda(x) := \nu_\lambda((-\infty, x])$ solves (3.9), and the special case $\lambda = \frac{1}{2}$ coincides with F above.

Let

$$W(x) = \cos^2\left(\frac{2\pi x}{\lambda}\right) \quad (x \in \mathbb{R}).$$

We saw in Example 3.2 that, for the IFS $\tau_0(x) = \lambda x$, $\tau_{1/4}(x) = \lambda(x + \frac{1}{4})$, the weight function W satisfies the normalization condition

$$R_W 1 = 1.$$

Also the attractor of this IFS is $X_{L(\lambda)} = [0, \frac{\lambda}{4(1-\lambda)}]$ if $\lambda > 1/2$. We are now ready to list the W -cycles for the Bernoulli convolutions for λ in the interval $(0, 1)$.

We will show that for the Bernoulli IFS systems with parameter λ , there are no W - p -cycles for $p > 1$, when λ is in the open interval $(0, 1)$.

Theorem 3.3. *Suppose $\lambda \in (0, 1)$. Then there are no W - p -cycles for $p > 1$.*

Proof. We relabel $\tau_1 := \tau_{1/4}$. Assume a priori that C is some W - p -cycle. Since it is not a W -1-cycle, the word ω which generates C must contain both the letters 0 and 1 from the alphabet $\{0, 1\}$.

First note that

$$(3.10) \quad W(x) = 1 \text{ only if } 2x/\lambda \in \mathbb{Z}.$$

Let x be now a point in a W - p -cycle, with $p > 1$. A p -cycle is generated by a word $\omega := (\omega_1 \dots \omega_p)$ of length p over the alphabet $\{0, 1\}$, i.e., $x = \pi_\lambda(\omega^\infty)$, where

$$\omega^\infty = \underbrace{(\omega \omega \omega \dots)}_{\text{infinite repetition}}.$$

But

$$\pi_\lambda(\omega^\infty) = \frac{1}{4} \lambda (\omega_1 + \omega_2 \lambda + \dots + \omega_p \lambda^{p-1}) \frac{1}{1 - \lambda^p}.$$

Therefore, with equation (3.10), we have that

$$(3.11) \quad (\omega_1 + \omega_2 \lambda + \dots + \omega_p \lambda^{p-1}) \frac{1}{2(1 - \lambda^p)} \in \mathbb{Z}.$$

We claim that λ must be rational. Indeed, after a cyclic permutation we may assume $\omega_1 = 0$. Then look at the next point in the cycle, which is generated by the

word $(\omega_2\omega_3\ldots\omega_p\omega_1)$. Clearly, this point will be λx . Using the fact that x and λx are points in a W -cycle, so $W(x) = W(\lambda x) = 1$, we obtain that

$$2x/\lambda \in \mathbb{Z}, \quad \text{and } 2x \in \mathbb{Z}.$$

This implies that λ is rational.

Let $\lambda = a/b$ with $a, b \in \mathbb{N}$, $a < b$ and a, b mutually prime. Using (3.11), we obtain that

$$b(\omega_1 b^{p-1} + \omega_2 b^{p-2}a + \cdots + \omega_p a^{p-1}) \frac{1}{2(b^p - a^p)} \in \mathbb{Z}.$$

If a prime number divides $b^p - a^p$ then it cannot divide b , because, if it does, it must also divide $b^p - (b^p - a^p) = a^p$. Then it would divide both a and b , which contradicts the fact that a and b are mutually prime.

Therefore we must have that $b^p - a^p$ divides $\omega_1 b^{p-1} + \omega_2 b^{p-2}a + \cdots + \omega_p a^{p-1}$. However

$$\omega_1 b^{p-1} + \omega_2 b^{p-2}a + \cdots + \omega_p a^{p-1} < b^{p-1} + b^{p-2}a + \cdots + a^{p-1} = \frac{b^p - a^p}{b - a} \leq b^p - a^p.$$

The inequality is strict, because at least one of the ω_i 's is 0. We have reached a contradiction, and the proof is complete. We conclude that when $\lambda \in (0, 1)$, there are no W -cycles of (minimal) period bigger than 1. $\square$

4. A HARMONIC FUNCTION

In this section, we address the problem of finding all the harmonic functions for the infinite Bernoulli convolutions defined for λ in the open interval $(0, 1)$. For every λ in $(0, 1)$, we are studying the corresponding Bernoulli-IFS. The present section has two main results; first a lemma (Lemma 4.1) about the zeroes of IFS-harmonic functions; and secondly our main result (Theorem 4.6): If λ is assumed irrational, then the constant functions are the only continuous IFS-harmonic functions. Specifically, the eigenspace H_λ in $C(X_\lambda)$ for the transfer operator R_{W_λ} with eigenvalue 1 is spanned by the constant function $h = 1$.

Recall that R_{W_λ} denotes the basic Ruelle operator (3.6), and $W = W_\lambda$ is defined in (3.4). And note further that we are addressing only the question of continuous solutions to $R_{W_\lambda}h = h$. In fact, it is intriguing that the case when λ is rational is the one that is more subtle as far as the IFS-harmonic functions are concerned. Finally, if measurable eigenfunctions are admitted into the analysis, then there are many more solutions; see, e.g., [Ba00], [Jor06], [JoPe98], [Nu01], and [LMW96]. We shall now apply the encoding $\mathbf{N}_C$ from Definition 2.5 of the possible W -cycles to the Bernoulli systems. We already saw that for the Bernoulli systems, the only possibilities for W - p -cycles are $p = 1$; and that then only C_0 and C_1 are possible. Which combinations actually occur depends on a partition of the value of λ (details below in Lemma 4.1).

Let

$$\mathbf{N}_0 := \{\omega = (\omega_1\omega_2\ldots) \in \{0, 1/4\}^\mathbb{N} \mid \omega_n = 0, \text{ for } n \text{ big enough}\},$$

i.e., $\mathbf{N}_0$ is the set of infinite paths that end in $000\ldots$.

We saw that if $\lambda = 1 - \frac{1}{2^n}$ for some $n \in \mathbb{N}$, then there is another W -cycle of length 1, namely $\{\frac{\lambda}{4(1-\lambda)}\}$. In this case we also define

$$\mathbf{N}_1 := \{\omega = (\omega_1\omega_2\ldots) \in \{0, 1/4\}^\mathbb{N} \mid \omega_n = 1/4, \text{ for } n \text{ big enough}\},$$

i.e., $\mathbf{N}_1$ is the set of infinite paths that end in $\frac{1}{4}\frac{1}{4}\frac{1}{4}\dots$.

Then let, for $x \in X_L$,

$$(4.1) \quad h_0(x) := P_x(\mathbf{N}_0) = \sum_{\omega \in \mathbf{N}_0} P_x(\{\omega\}) = \sum_{\omega \in \mathbf{N}_0} \prod_{k=1}^{\infty} W(\tau_{\omega_k} \cdots \tau_{\omega_1} x).$$

In the case when $\lambda = 1 - \frac{1}{2^n}$, we also define

$$(4.2) \quad h_1(x) := P_x(\mathbf{N}_1) = \sum_{\omega \in \mathbf{N}_1} P_x(\{\omega\}) = \sum_{\omega \in \mathbf{N}_1} \prod_{k=1}^{\infty} W(\tau_{\omega_k} \cdots \tau_{\omega_1} x).$$

Lemma 4.1. *Consider the functions h_0 and h_1 defined above.*

- (i) *The function h_0 is continuous and harmonic for the transfer operator R_W , i.e., $R_W h_0 = h_0$. If h_0 has no zeroes then $h_0 = 1$.*
- (ii) *If $\lambda = 1 - \frac{1}{2^n}$ then h_1 is continuous and harmonic for the transfer operator R_W . Both functions then have zeroes, and both are non-constant. If $h_0 + h_1$ has no zeroes then $h_0 + h_1 = 1$.*

Proof. We shall use the one-to-one correspondence from Definition 2.5, between cycles C and harmonic functions h_C ; and we first consider the cycle $C_0 = \{0\}$, setting $h_0 := h_{C_0}$. The fact that h_0 is continuous and harmonic follows from [DuJo05]. Also, from [DuJo05, Theorem 4.1] we know that

$$\chi_{\mathbf{N}_0}(\omega) = \lim_{k \rightarrow \infty} h_0(\tau_{\omega_k} \cdots \tau_{\omega_1} x), \quad \text{for } P_x \text{ a.e. } \omega.$$

So, if h_0 has no zeroes then $\mathbf{N}_0$ must have P_x -measure 1, otherwise there are some ω for which the limit is 0, and, using the continuity of h_0 , we obtain a contradiction. The argument in the proof of the second assertion in the lemma is essentially the same, with the modification that we now consider the union of the two subsets $\mathbf{N}_0$ and $\mathbf{N}_1$. Assuming that $h_0 + h_1$ has no zeroes, then the same reasoning as above yields $\chi_{\mathbf{N}_0 \cup \mathbf{N}_1} = 1$ P_x a.e., and therefore $P_x(\mathbf{N}_0 \cup \mathbf{N}_1) = 1 = h_0(x) + h_1(x)$. $\square$

Remark 4.2. Our limit formula above for the indicator function $v := \chi_{\mathbf{N}_0}$ is a special case of a martingale limit argument from [DuJo05, Theorem 3.1], see also [Gun00]. Our result from [DuJo05] is further a close analogue for IFS-harmonic functions of the familiar and classical Fatou-Primalov boundary-value theorem in Fourier analysis [Rud87].

A special class of cocycles v in [DuJo05] are the indicator functions of subsets $\mathbf{N}_C$ of the symbol space Ω which are defined in terms of the shift on Ω ; i.e., $v = \chi_{\mathbf{N}_C}$. The reader is referred to [DuJo05] for additional details and to the list of cycles for the Bernoulli convolutions above, i.e., the list before the statement of Theorem 3.3.

We first need a definition and some technical results.

Definition 4.3. Let $\lambda \in \mathbb{R}_+$, and suppose for some $p \geq 2$, $\lambda^p \in \mathbb{Q}$. We say that p is minimal if $\lambda^{p_1} \notin \mathbb{Q}$ when $0 < p_1 < p$. The number p is called the minimal degree of λ .

The minimal polynomial $m(x) \in \mathbb{Q}[x]$ of λ is a monic polynomial, $m(\lambda) = 0$, of lowest degree with λ as root. It is irreducible in $\mathbb{Q}[x]$; see [Hun84, p. 234]. Moreover, if $f(x) \in \mathbb{Q}[x]$ satisfies $f(\lambda) = 0$, then $m(x)$ divides $f(x)$.

Lemma 4.4. *If $\lambda = q^{\frac{1}{p}}$ with $q \in \mathbb{Q}$ and $p \geq 2$ minimal, then the minimal polynomial of λ is $x^p - q$.*

Proof. Let $m(x)$ be the minimal polynomial of λ . Then $m(x)$ has to divide $x^p - q$, therefore $m(x)$ has to be of the form

$$m(x) = \prod_k (x - \lambda z_k),$$

where the product is over some of the p 'th roots of unity. Let $1 \leq l \leq p$ be the degree of m . The constant term is of the form $\lambda^l z$ for some $z \in \mathbb{C}$ with $|z| = 1$. At the same time it has to be rational. Thus $z = \pm 1$ and λ^l is rational. Since p is minimal, it follows that $p = l$ and that $m(x) = x^p - q$. $\square$

Corollary 4.5. *Let $\lambda \in \mathbb{R}_+$, and let p be its minimal degree, $p \geq 2$. Then there are no solutions*

$$\lambda^r = a\lambda^s + b,$$

with $a, b \in \mathbb{Q}$, and $0 \leq s < r < p$.

Proof. If λ satisfies this condition, then the polynomial $x^r - ax^s - b$ would be divisible by $x^p - q$; and this is impossible since $r < p$. $\square$

Theorem 4.6. *Let λ be in $(0, 1)$.*

- (i) *If $\lambda \notin \{1 - \frac{1}{2^n} \mid n \in \mathbb{N}\}$ then $h_0 = 1$.*
- (ii) *If $\lambda = 1 - \frac{1}{2^n}$ then $h_0 + h_1 = 1$.*

Proof. We use Lemma 4.1, and prove that h_0 has no zeroes. Using (4.1), we see that it is enough to prove that, for each $x \in X_L$, there is a path ω that ends in $000\dots$ (or in $\frac{1}{4}\frac{1}{4}\frac{1}{4}\dots$ for (ii)) which does not go through zeroes of W , i.e., $\tau_k \cdots \tau_1 x$ is not a zero of W for all $k \geq 1$.

Since $W = \cos^2\left(\frac{2\pi x}{\lambda}\right)$, the zeroes of W are $Z := \{\frac{\lambda}{2}(\frac{1}{2} + n) \mid n \in \{0, 1, \dots\}\}$. Define now the set

$$A := \bigcup_{p \geq 1} \lambda^{-p} Z.$$

If $x \notin A$ then the desired path is $\omega = 000\dots$.

Assume now that $x \in A$. Take the smallest $p \geq 1$, such that $\lambda^p x \in Z$. There exists $m \in \{0, 1, \dots\}$ and $p \geq 1$ such that $x = \lambda^{-p} \frac{\lambda}{2}(\frac{1}{2} + m)$. Since p is minimal, $\tau_0^k x \notin Z$ for $k \leq p-1$.

We consider first the case when $\lambda \in (\frac{1}{2}, 1)$ is not of the form $\lambda = \left(\frac{n+1/2}{m+1}\right)^{\frac{1}{p}}$, for $m, n \in \{0, 1, \dots\}$, $p \geq 1$.

We prove now that $\tau_{1/4} \tau_0^{p-1} x \notin A$. If not, then there exist $r \geq 1$ and $n \in \{0, 1, \dots\}$ such that

$$\lambda \left(\lambda^{-1} \frac{\lambda}{2} \left(\frac{1}{2} + m \right) + \frac{1}{4} \right) = \lambda^{-r} \frac{\lambda}{2} \left(\frac{1}{2} + n \right).$$

This implies that $\lambda^r = \frac{n+1/2}{m+1}$, which contradicts the hypothesis.

Thus, in this case, the desired path is $\underbrace{00\dots 0}_{p-1 \text{ times}} \frac{1}{4} 00\dots$.

We can assume now that λ is of the form $\lambda = q^{\frac{1}{p}}$ for some q rational and $p \geq 1$.

The argument above shows that we may assume that $x \in \lambda^{-1} Z$, $x = \lambda^{-1} \frac{\lambda}{2}(n + \frac{1}{2}) = \frac{n}{2} + \frac{1}{4}$, for some natural number $n \geq 0$. Then $W(\tau_0 x) = 0$ so $W(\tau_{1/4} x) = 1$. We prove that $\tau_{1/4} \tau_{1/4} x \notin A \cup Z$.

Suppose $\tau_{1/4}\tau_{1/4}x$ is in $A \cup Z$, then for some $r \geq 0$ and $m \in \{0, 1, \dots\}$ we have

$$\lambda \left(\lambda \left(\left(\frac{n}{2} + \frac{1}{4} \right) + \frac{1}{4} \right) + \frac{1}{4} \right) = \lambda^{-r} \frac{\lambda}{2} \left(m + \frac{1}{2} \right),$$

so

$$(4.3) \quad \lambda^{r+1} \left(\frac{n}{2} + \frac{1}{2} \right) + \lambda^r \frac{1}{4} = \left(\frac{m}{2} + \frac{1}{4} \right).$$

We will show that this yields a contradiction with the fact that λ^p is rational for some $p \geq 2$.

We first need a definition and some technical results.

Consider p minimal such that $\lambda^p \in \mathbb{Q}$. If in equation (4.3), $r+1 \geq p$ then we can write $r+1 = pi + j$ for $i \in \mathbb{N}$ and $j < p$. Then $\lambda^{r+1} = \lambda^{pi} \lambda^j \in \mathbb{Q} \lambda^j$, therefore, using (4.3), we see that λ satisfies an equation of the form $\lambda^r = a\lambda^s + b$ with $0 \leq s < r < p$, and $a, b \in \mathbb{Q}$, and this contradicts Corollary 4.5.

Finally we obtain that $\frac{1}{4}\frac{1}{4}000\dots$ is the desired path that avoids zeroes of W .

The remaining case is when λ is rational. Take $x \in X_L$. If $x = 0$ then the path is $000\dots$. In (ii) if $x = \frac{\lambda}{4(1-\lambda)}$, then the path is $\frac{1}{4}\frac{1}{4}\frac{1}{4}\dots$.

Let $x \in X_L \setminus \{0, \frac{\lambda}{4(1-\lambda)}\}$. We say that $(\omega_1, \dots, \omega_n)$ is a path that starts at x and avoids zeroes if $W(\tau_{\omega_1}x), \dots, W(\tau_{\omega_n} \cdots \tau_{\omega_1}x)$ are all non-zero. Let r_n be the number of paths that avoid zeroes. Then $r_{n+1} \geq r_n$ because for any path $(\omega_1, \dots, \omega_n)$ that avoids zeroes, since

$$W(\tau_0(\tau_{\omega_n} \cdots \tau_{\omega_1}x)) + W(\tau_{1/4}(\tau_{\omega_n} \cdots \tau_{\omega_1}x)) = 1,$$

there is a $\omega_{n+1} \in \{0, 1/4\}$ such that $(\omega_1, \dots, \omega_{n+1})$ is a path that avoids zeroes.

Assume that r_n is bounded. Then for some n_0 , $r_{n+1} = r_n$ for all $n \geq n_0$. Denote by

$$E_n := \{\tau_{\omega_n} \cdots \tau_{\omega_1}x \mid (\omega_1, \dots, \omega_n) \text{ is a path that avoids zeroes}\}.$$

Since $r_n = r_{n+1}$ for $n \geq n_0$, it means that for every $y \in E_n$, $n \geq n_0$, there is only one choice to continue the path that starts at x and ends at y and avoids zeroes (otherwise $r_{n+1} \geq r_n + 1$). Thus $W(\tau_0y) = 0$ or $W(\tau_{1/4}y) = 0$.

This implies that $y \in \tau_0^{-1}(Z) \cup \tau_{1/4}^{-1}(Z)$ where Z is the set of zeroes of W . Consequently $E := \bigcup_{n \geq n_0} E_n$ is a finite set.

Now pick y in E . We have that τ_0x or $\tau_{1/4}x$ is a zero for W . Therefore $W(\tau_{1/4}x) = 1$ or $W(\tau_0x) = 1$. Let $\eta_1 \in \{0, 1/4\}$ such that $W(\tau_{\eta_1}y) = 1$. Then $\tau_{\eta_1}y \in E$. Inductively we can construct $\eta_1, \dots, \eta_m$ such that $W(\tau_{\eta_k} \cdots \tau_{\eta_1}x) = 1$ and $\tau_{\eta_k} \cdots \tau_{\eta_1}x \in E$. Since E is finite, $\tau_{\eta_k} \cdots \tau_{\eta_1}x$ must end in a cycle. Moreover, the previous argument shows that this will be a W -cycle.

However the only cycles are the 1-cycles 0 and possibly $\frac{\lambda}{4(1-\lambda)}$ in (ii). But note that for $z \in X_L$, $\tau_\alpha z = 0$ implies $z = 0$, and $\tau_\alpha z = \frac{\lambda}{4(1-\lambda)}$ implies $z = \frac{\lambda}{4(1-\lambda)}$. Thus $x \in \{0, \frac{\lambda}{4(1-\lambda)}\}$.

Thus we have that r_n is unbounded. Next we prove that, since λ is rational, we cannot have $\tau_{\omega_n} \cdots \tau_{\omega_1}x = \tau_{\eta_n} \cdots \tau_{\eta_1}x$ for $(\omega_1, \dots, \omega_n) \neq (\eta_1, \dots, \eta_n)$. Indeed, if they are equal then we obtain

$$\omega_n + \lambda\omega_{n-1} + \cdots + \lambda^{n-1}\omega_1 = \eta_n + \lambda\eta_{n-1} + \cdots + \lambda^{n-1}\eta_1.$$

Then λ will satisfy an equation of the form

$$a_r \lambda^r + \cdots + a_1 \lambda + a_0 = 0,$$

where $a_i \in \{0, 1, -1\}$, not all of them 0, and $r \geq 1$. Let $\lambda = a/b$ with a, b mutually prime integers. We have

$$a_r a^r + a_{r-1} a^{r-1} b + \cdots + a_0 b^r = 0.$$

But all the terms are divisible by b except the first one, and this yields a contradiction.

Since r_n is unbounded and, since for different paths of the same length starting at x the endpoints are different, it follows that the cardinality of the set E_n increases to infinity.

However, note that the set A is finite because Z is finite and $\lambda^{-r} z \notin X_L$ for r large. It follows then, that if we choose n such that the cardinality of E_n is bigger than the one of A , there is a path of length n that starts at x , avoids zeroes, and ends outside A . Hence this path continued with $000 \dots$ is the desired one. $\square$

Corollary 4.7. *Consider the Bernoulli IFSs for all λ in the open interval $(0, 1)$. For λ fixed, and for all x , each of the measures P_x on Ω is countably atomic. By this we mean that each P_x is supported on a certain countable subset S of Ω , and each point ω in S is an atom for P_x .*

Remark 4.8. This property from the conclusion of the corollary for the process P_x is called “tightness”. See, e.g., [Gun00], [Jor06], and [Fal93], where an analogous process (Ω, P_x) is studied in a different context: orthogonality relations for the wavelet scaling function. Our result, Corollary 4.7 for the Bernoulli convolutions, to the effect that the measures P_x are purely atomic, stands in contrast with a different result below, for the Bernoulli measures ν_λ . In fact, we show in Corollary 6.6 that the strongly invariant measures for affine IFSs (which includes the Bernoulli measures ν_λ) are all non-atomic.

Let $\lambda \in (0, 1)$ and consider the random variable $\sum_{j=0}^{\infty} (\pm 1) \lambda^j$, i.e., independently distributed ones and minus ones. It follows from the previous sections that the distribution of this random variable is the measure $\nu_\lambda = p_{(\frac{1}{2}, \frac{1}{2})} \circ \pi_\lambda^{-1}$.

Lemma 4.9. *Let $\lambda \in (0, 1)$, and let ν_λ be the corresponding infinite convolution IFS measure. Then the Fourier transform*

$$(4.4) \quad \hat{\nu}_\lambda(t) := \int e^{2\pi i x t} d\nu_\lambda(x)$$

satisfies

$$(4.5) \quad \hat{\nu}_\lambda(t) = \prod_{n=0}^{\infty} \cos(2\pi \lambda^n t).$$

Proof. By Lemma 6.1, (6.3), we get

$$(4.6) \quad \nu_\lambda = \frac{1}{2} (\nu_\lambda \circ \tau_+^{-1} + \nu_\lambda \circ \tau_-^{-1}),$$

or equivalently,

$$\int f(x) d\nu_\lambda(x) = \frac{1}{2} \left(\int f(\tau_+(x)) d\nu_\lambda(x) + \int f(\tau_-(x)) d\nu_\lambda(x) \right),$$

see (3.2). Using this and (4.4), we get

$$\begin{aligned}
\hat{\nu}_\lambda(t) &= \int e^{2\pi i x t} d\nu_\lambda(x) \\
&= \frac{1}{2} \left(\int e^{2\pi i(\lambda x + 1)t} d\nu_\lambda(x) + \int e^{2\pi i(\lambda x - 1)t} d\nu_\lambda(x) \right) \\
&= \frac{e^{2\pi i t} + e^{-2\pi i t}}{2} \int e^{2\pi i \lambda x t} d\nu_\lambda(x) = \cos(2\pi t) \hat{\nu}_\lambda(\lambda t) \\
&= \cdots = \prod_{k=0}^{n-1} \cos(2\pi \lambda^k t) \hat{\nu}_\lambda(\lambda^n t),
\end{aligned}$$

by iteration.

A limit argument, using that

$$\lim_{n \rightarrow \infty} \hat{\nu}_\lambda(\lambda^n t) = \hat{\nu}_\lambda(0) = 1,$$

now yields the desired infinite-product formula (4.5). $\square$

We will use the following notation: for a finite word $\omega = \omega_1 \dots \omega_n \in \{0, 1/4\}^n$, we denote by W^ω and τ_ω the following objects:

$$W^\omega(x) := W(\tau_{\omega_n} \cdots \tau_{\omega_1} x) \cdots W(\tau_{\omega_1} x), \quad W^\emptyset(x) = 1,$$

$$\tau_\omega x = \tau_{\omega_n} \cdots \tau_{\omega_1} x, \quad \tau_\emptyset x = x.$$

Theorem 4.10. *Let $\lambda \in (0, 1)$, λ not of the form $\lambda = 1 - \frac{1}{2^n}$, $n \in \mathbb{N}$. Let*

$$\mathcal{W}_0 := \{\emptyset\} \cup \{\omega = \omega_1 \dots \omega_n \mid \omega_i \in \{0, 1/4\}, \omega_n = 1/4, n \geq 1\}.$$

The function $|\hat{\nu}_\lambda|^2$ is the only continuous function of X_λ that satisfies the following identity

$$(4.7) \quad \sum_{\omega \in \mathcal{W}_0} W^\omega(x) f(\tau_\omega x) = 1, \quad (x \in X_L).$$

Proof. First we prove that $|\nu_\lambda|^2$ satisfies (4.7). From Theorem 4.6 we know that

$$(4.8) \quad P_x(\mathbf{N}_0) = 1.$$

Take $\omega \in \mathbf{N}_0$. If $\omega = 000 \dots$, then, from [DuJo05], we know that

$$P_x(\{000 \dots\}) = \prod_{k=1}^{\infty} W(\tau_0 \cdots \tau_0 x) = |\hat{\nu}_\lambda(x)|^2,$$

where we used Lemma 4.9 in the last equality.

If $\omega \neq 000 \dots$, then $\omega = \omega_1 \dots \omega_n 00 \dots$, where $\omega_n = 1/4$. Then

$$\begin{aligned}
P_x(\{\omega\}) &= W(\tau_{\omega_1} x) \cdots W(\tau_{\omega_n} \cdots \tau_{\omega_1} x) \prod_{k=1}^{\infty} W(\tau_0^k \tau_{\omega_n} \cdots \tau_{\omega_1} x) \\
&= W^{\omega_1 \dots \omega_n}(x) |\hat{\nu}_\lambda(\tau_\omega x)|^2.
\end{aligned}$$

Thus equation (4.8) can be rewritten as (4.7).

Take now a continuous function f on $X_L = [0, \frac{\lambda}{4(1-\lambda)}]$ which satisfies (4.7). We rewrite this identity, by enumerating the finite words that end in 1:

$$\begin{aligned} 1 &= f(x) + W(\tau_{1/4}x)f(\tau_{1/4}x) + \cdots \\ &\quad + \sum_{\omega_1, \dots, \omega_{n-1}} W(\tau_{\omega_0}x) \cdots W(\tau_{\omega_{n-1}} \cdots \tau_{\omega_1}x) \\ &\quad \cdot W(\tau_{1/4}\tau_{\omega_{n-1}} \cdots \tau_{\omega_1}x)f(\tau_{1/4}\tau_{\omega_{n-1}} \cdots \tau_{\omega_1}x) + \cdots . \end{aligned}$$

Therefore

$$(4.9) \quad 1 = f(x) + R_W^0(W \circ \tau_{1/4} f \circ \tau_{1/4})(x) + \cdots + R_W^{n-1}(W \circ \tau_{1/4} f \circ \tau_{1/4})(x) + \cdots .$$

In particular, $R_W^n(W \circ \tau_{1/4} f \circ \tau_{1/4})$ converges pointwise to 0.

We apply the operator $I - R_W$ to (4.9). (More precisely, we consider equation (4.9) at $\tau_0 x$ and $\tau_{1/4} x$, and multiply by $W(\tau_0 x)$ and $W(\tau_{1/4} x)$ respectively. Then we subtract both these equations from (4.9). All the series involved in the computation converge pointwise). We obtain

$$\begin{aligned} 0 &= f - R_W f + W \circ \tau_{1/4} f \circ \tau_{1/4} - R_W(W \circ \tau_{1/4} f \circ \tau_{1/4}) + \cdots \\ &\quad + R_W^{n-1}(W \circ \tau_{1/4} f \circ \tau_{1/4}) - R_W^n(W \circ \tau_{1/4} f \circ \tau_{1/4}) + \cdots \\ &= f - R_W f + W \circ \tau_{1/4} f \circ \tau_{1/4} = f - W \circ \tau_0 f \circ \tau_0 . \end{aligned}$$

Thus f satisfies the equation

$$(4.10) \quad f(x) = W(\lambda x)f(\lambda x), \quad x \in \left[0, \frac{\lambda}{4(1-\lambda)}\right] .$$

Note also that equation (4.9) implies that $f(0) = 1$. Then, using the relations in the proof of Lemma 4.9, we see that the function $g := f/|\hat{\nu}_\lambda|^2$ verifies the equation

$$g(x) = g(\lambda x),$$

for all x in $[0, \frac{\lambda}{4(1-\lambda)}]$ except when x is a zero for $\hat{\nu}_\lambda$. The zeroes of $\hat{\nu}_\lambda$ inside the interval X_L are finite in number, because they are of the form $\lambda^{-k}z$ where z is a zero of W .

In addition, the function g is continuous outside this finite set; it is also continuous at 0 and $g(0) = 1$. This implies that $g(x) = g(\lambda^n x) = \lim_{n \rightarrow \infty} g(\lambda^n x) = g(0) = 1$ for x outside some finite set. Finally we can conclude that $g = 1$ except at some finite number of points, and this implies that $f = |\hat{\nu}_\lambda|^2$. $\square$

Proposition 4.11. *The convergence in (4.7) is uniform on $X_\lambda = [0, \frac{\lambda}{4(1-\lambda)}]$.*

Proof. By Theorem 4.10, we have that $f = |\hat{\nu}_\lambda|^2$. But this is a non-negative continuous function and the same is true for W . Thus the uniform convergence follows from the pointwise convergence and Dini's theorem. $\square$

5. λ REPRESENTATIONS

While there is already a substantial literature on the structure of infinite Bernoulli convolutions, we take here a different approach. In this paper, the systems (X_λ, ν_λ) serve as a link between the class of iterated function systems (IFS) and the more general scale-similarity notions in analysis, e.g., the scale covariance which is intrinsic to wavelet theory; see the discussion around equations (3.8) and (3.9) above. We showed in [DuJo05] that in general IFSs carry a discrete harmonic analysis which is based on W -cycles. For general IFSs, the family of W -cycles may in fact be

quite complicated and not especially computable. Our present Theorem 3.3 in the previous section shows that for each of the Bernoulli systems X_λ , we can have at most two W -cycles. We found them, and worked out the harmonic analysis. This harmonic analysis as a result is somewhat coarse.

The purpose of this section is to relate this theorem and its implications to the geometry of the Bernoulli systems X_λ and to their encoding mappings $\pi_\lambda: \Omega \rightarrow X_\lambda$. While it is possible that some of our results spelled out in Proposition 5.1 below are known, we have not been able to locate them in the literature.

For a parameter λ in the open interval $(0, 1)$, consider the random variable $\sum_k (\pm \lambda^k)$, with independent $\frac{1}{2}$ - $\frac{1}{2}$ -distribution of the $\pm$ signs in the series, and let ν_λ be the corresponding distribution measure. Solomyak [So95] showed that for Lebesgue a.e. λ in $(1/2, 1)$, ν_λ has an L^2 density. Nonetheless, when $\lambda > 1/2$, precious little is known about the explicit properties of this one-parameter family of measures ν_λ . A fundamental result is due to Paul Erdős [Er39] who showed that ν_λ is purely singular when the reciprocal λ^{-1} is a Pisot number, i.e., an algebraic number whose Galois conjugates are strictly less than one in absolute value. While P. Erdős' theorem [Er39] stands out, it has now been extended to a few larger classes of algebraic numbers (see especially the deep results by Garcia [Ga62] about absolute continuity, and Kahane [Kah71] proving that the Fourier transform of ν_λ does not vanish at infinity when λ^{-1} is a Salem number), yet there is little known about specific values of λ ; for example, few explicit properties are known about the measure $\nu_{2/3}$. Yet, the isolated results in the literature do depend on certain algebraic/arithmetic properties of the number λ . In fact the Diophantine properties of λ will play a role in our present analysis.

In order to apply our Fourier duality to the classical λ -systems, we need a few geometric facts about these systems. We have therefore assembled them in the next proposition. While they may be known, we have not been able to find them exactly in the form that is needed for our present harmonic analysis. So we include them without proof for the benefit of the reader.

Proposition 5.1. *Let $\lambda \in (0, 1)$ and $B = \{0, 1\}$, and let (τ_b) , X_B be the corresponding affine IFS.*

- (a) *If $\lambda < 1/2$, then X_B is a fractal.*
- (b) *If $\lambda \geq 1/2$, then*

$$(5.1) \quad X_B = \left[0, \frac{\lambda}{1-\lambda} \right],$$

i.e., a closed interval.

- (c) *If $0 < \lambda < 1/2$, then $\pi_\lambda: \Omega \rightarrow X_B$ is a bijection.*
- (d) *If $\frac{1}{2} < \lambda < 1$, then for every point $x \in (0, \frac{\lambda}{1-\lambda})$, $\pi_\lambda^{-1}(x)$ is uncountable.*

Proof. See [JoPe96], [Edg98], [JeWi35], [KeWi35], [Anll01] and the references therein. $\square$

Corollary 5.2. *If $0 < \lambda < \frac{1}{2}$, then X_B is a Cantor set of Lebesgue measure zero and Hausdorff dimension $d_H = \frac{\ln(2)}{\ln(\lambda^{-1})}$.*

Proof. We first show that the Lebesgue measure of the support set X_λ is a fractal, and that its Lebesgue measure is zero. Start with the interval $[0, \frac{\lambda}{1-\lambda}]$ and omit the

middle open interval $(\frac{\lambda^2}{1-\lambda}, \lambda)$ of length $\frac{\lambda(1-2\lambda)}{1-\lambda}$. Continue to omit the middle open intervals at each λ -scale.

The sum of all the omitted middle fractions is

$$\frac{\lambda(1-2\lambda)}{1-\lambda}(1+2\lambda+(2\lambda)^2+\dots) = \frac{\lambda(1-2\lambda)}{1-\lambda} \frac{1}{1-2\lambda} = \frac{\lambda}{1-\lambda},$$

so the omitted intervals have full measure in the interval $[0, \frac{\lambda}{1-\lambda}]$. The stated formula for the Hausdorff dimension d_H now follows from the discussion above, and from standard scaling theory; see, e.g., [Edg98]. $\square$

Remark 5.3. For $y \in \mathbb{R}$, set $\langle y \rangle := y - \lfloor y \rfloor$ = the fractional part of y . Then $0 \leq \langle y \rangle < 1$. When $0 < \lambda < 1$ is given, set $r_\lambda: [0, 1) \rightarrow [0, 1)$,

$$(5.2) \quad r_\lambda(y) := \langle \lambda^{-1}y \rangle$$

We note that this is a dynamical system. This system has uses in symbolic dynamics, for example in the study of subshifts, see [Wal78].

We now recall a recursive number-representation based on (5.2). It is simply a recursive algorithm for a λ -representation of certain real numbers x . When $\lambda < 1/2$, it does yield the representation of points in the fractal X_λ , but if $\lambda \geq 1/2$ (Lemma 5.4), this representation typically has two defects: first, the required alphabet A is then bigger than $\{0, 1\}$; and secondly, the dynamical systems expansion (5.3) only offers *one* representation of the admissible numbers x . Our result Proposition 5.1 above in fact keeps track of the number of different representations for the same x ; and the alphabet A in Proposition 5.1 is fixed, $A = \{0, 1\}$.

When $0 < \lambda < \frac{1}{2}$, we saw that the expansion $x = \sum_{k=1}^{\infty} \omega_k \lambda^k$, is unique, $\omega_k \in \{0, 1\}$ for all $x \in X_B$. An application of (5.2) shows that

$$(5.3) \quad \omega_k = \left\lfloor \frac{1}{\lambda} r_\lambda^{k-1}(x) \right\rfloor.$$

If $0 < \lambda < \frac{1}{2}$, then there are only two possibilities, 0 or 1, for ω_k . Not so if $\frac{1}{2} \leq \lambda < 1$.

There is a more general expansion $x = \sum_{i=1}^{\infty} \omega_i \lambda^i$ which has the π_λ expansion as a special case when $\lambda < \frac{1}{2}$.

For $y \in \mathbb{R}$, let $\lfloor y \rfloor$ = integral part, $\langle y \rangle := y - \lfloor y \rfloor$ = fractional part.

If $0 < \lambda < 1$, set a = the largest integer with $a < \frac{1}{1-\lambda}$. Set

$$r_\lambda(y) := \left\langle \frac{1}{\lambda} y \right\rangle, \quad \omega_k := \left\lfloor \frac{1}{\lambda} r_\lambda^{k-1}(x) \right\rfloor \in \{0, 1, \dots, a\}.$$

Lemma 5.4. *Let $0 < \lambda < 1$, and let A be the set of integers $\{0, 1, \dots, a\}$ where $a < \frac{1}{1-\lambda}$. (Note that if $\lambda < \frac{1}{2}$, then $A = \{0, 1\}$.) Then every $x \in X_B$ has a convergent expansion*

$$(5.4) \quad x = \sum_{k=1}^{\infty} \omega_k \lambda^k,$$

with $\omega_k \in A$ as in (5.3).

Proof. We only need to show that (5.4) is convergent. But the recursive algorithm shows that

$$0 \leq x - \sum_{j=1}^k \omega_j \lambda^j \leq a \frac{\lambda^k}{1-\lambda},$$

where a is the largest integer $< \frac{1}{1-\lambda}$. Since $\lambda < 1$, the result follows. $\square$

The following simple lemma shows that the IFS associated to $\lambda \in (0, 1)$ and an arbitrary pair $B = \{a, b\}$ is equivalent up to some scaling and translation to the one associated to $\{0, 1\}$. Therefore in analyzing these systems, we can take any pair B .

Lemma 5.5. *Let $\lambda \in (0, 1)$ and $B = \{a, b\}$ with $a < b$ in $\mathbb{R}$. Then the affine function $\alpha(x) = (b-a)x + a\frac{\lambda}{1-\lambda}$ has the property that, for $(a_i)_i \in \{0, 1\}^{\mathbb{N}}$,*

$$\alpha\left(\sum_{i=1}^{\infty} a_i \lambda^i\right) = \sum_{i=1}^{\infty} b_i \lambda^i, \text{ where } b_i := \begin{cases} a & \text{if } a_i = 0, \\ b & \text{if } a_i = 1. \end{cases}$$

Proof. It is enough to see that

$$a \sum_{i=1}^{\infty} a_i \lambda^i + (b-a) \frac{\lambda}{1-\lambda} = \sum_{i=1}^{\infty} (aa_i + (b-a)) \lambda^i.$$

$\square$

6. STRONGLY INVARIANT MEASURES

Let $(X, (\tau_i)_{i=1}^N)$ be an IFS, and suppose it has an encoding mapping $\pi: \Omega \rightarrow X$ which is measurable with respect to two σ -algebras

- $\mathfrak{B}(\Omega)$: the σ -algebra on Ω which is generated by the cylinder sets.
- $\mathfrak{B}(X)$: the Borel σ -algebra on X .

For every $(p_i)_{i=1}^N$, $p_i > 0$, $\sum_{i=1}^N p_i = 1$, there is a unique probability measure $\mu_{(p)}$, on Ω , called the Bernoulli measure, and determined by

$$(6.1) \quad \mu_{(p)}(\omega_1 \omega_2 \dots \omega_n \times \mathbb{Z}_N \times \mathbb{Z}_N \times \dots) = p_{\omega_1} p_{\omega_2} \dots p_{\omega_n}.$$

Lemma 6.1. *Let $(X, (\tau_i))$ be an IFS which has a measurable encoding mapping π . Let (p_i) be as above, and let $\mu_{(p)}$ be the Bernoulli measure on Ω . Then the measure*

$$(6.2) \quad \nu_{(p)}^X := \mu_{(p)} \circ \pi^{-1}$$

on X satisfies

$$(6.3) \quad \nu_{(p)}^X = \sum_{i=1}^N p_i \nu_{(p)}^X \circ \tau_i^{-1},$$

and

$$\text{supp}(\nu_{(p)}^X) = X,$$

where supp denotes the support.

Proof. The measure in (6.2) is defined by $\mu_{(p)}(\pi^{-1}(E))$ for $E \in \mathfrak{B}(X)$, where

$$\pi^{-1}(E) = \{\omega \in \Omega \mid \pi(\omega) \in E\};$$

and similarly

$$(\nu_{(p)}^X \circ \tau_i^{-1})(E) = \nu_{(p)}^X(\tau_i^{-1}(E)).$$

It follows from the definition of $\mu_{(p)}$ in (6.1) that

$$\mu_{(p)} = \sum_{i=1}^N p_i \mu_{(p)} \circ \sigma_i^{-1},$$

where $\sigma_i: \Omega \rightarrow \Omega$ is defined by $\sigma_i(\omega_1 \omega_2 \dots) := (i \omega_1 \omega_2 \dots)$. From (2.8), we conclude that

$$\pi \circ \sigma_i = \tau_i \circ \pi.$$

Let $E \in \mathfrak{B}(X)$. Then

$$\begin{aligned} \nu_{(p)}^X(E) &= \mu_{(p)}(\pi^{-1}(E)) = \sum_{i=1}^N p_i \mu_{(p)}(\sigma_i^{-1} \pi^{-1}(E)) \\ &= \sum_{i=1}^N p_i \mu_{(p)}(\pi^{-1} \tau_i^{-1}(E)) = \sum_{i=1}^N p_i \nu_{(p)}^X(\tau_i^{-1}(E)), \end{aligned}$$

which is the desired formula (6.3).

We now prove that $\nu_{(p)}^X$ has full support. The proof is indirect: Suppose $\emptyset \neq G \subset X$ is open and $\nu_{(p)}^X(G) = 0$. Then there is a finite word $(i_1 \dots i_n)$ such that

$$\sigma_{i_1} \cdots \sigma_{i_n}(\Omega) \subset \pi^{-1}(G).$$

But then

$$\nu_{(p)}^X(G) = \mu_{(p)}(\pi^{-1}(G)) \geq \mu_{(p)}(\sigma_{i_1} \cdots \sigma_{i_n} \Omega) = p_{i_1} \cdots p_{i_n},$$

contradicting that $p_i > 0$ for all i . Hence $\nu_{(p)}^X$ does not vanish on non-empty open subsets in X , which is to say

$$\text{supp}(\nu_{(p)}^X) = X;$$

see [Rud87, page 58]. □

Definition 6.2. A probability measure $\nu_{(p)}^X$ on X satisfying equation (6.3) is called strongly invariant.

Lemma 6.1 proves in particular that strongly invariant measures exist. When the maps τ_i are contractions the strongly invariant measure is also unique (see [Hut81]).

We show that a class of these strongly invariant measures are non-atomic, i.e., that

$$\nu_{(p)}^X(\{x\}) = 0, \quad \text{for } x \in X.$$

Definition 6.3. Let $(X, (\tau_i)_{i=1, \dots, N})$ be an IFS, and let x be in X . Then the backward orbit of x , $O^-(x)$ is defined as

$$O^-(x) := \bigcup \{ \tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1} x \mid n \in \mathbb{N}, \omega_1, \dots, \omega_n \in \{1, \dots, N\} \}.$$

Our result in this section is based on an *a priori* observation about atoms for strongly invariant measures ν (unique or not!) for IFSs. In this generality, our result states that if an IFS with one-to-one maps has atoms, it has a maximal atom, i.e., an atom x such that $a := \nu(\{x\}) \geq \nu(\{y\})$ for all other atoms y . We then show that all the non-empty sets in the backward orbit $O^-(x)$ from a maximal atom x consist of atoms of the same measure. We note that this fact strongly restricts those IFS that can have atoms; and we prove in Corollary 6.6 that all contractive affine IFS are non-atomic.

Theorem 6.4. *Let $(X, (\tau_i)_{i \in \{1, \dots, N\}})$ be an iterated function system, $(p_i)_{i=1}^N$ given as above, with $p_i > 0$ for all i , and let $\nu := \nu_{(p)}$ be the unique strongly invariant probability measure. Assume that for all $x \in X$, either the backward orbit of x is infinite, or there are some $\omega_1, \dots, \omega_n$ such that $\tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1}(x) = \emptyset$. Then the measure ν has no atoms.*

Proof. Since ν is strongly invariant, it satisfies the equation

$$(6.4) \quad \nu(\{x\}) = \sum_{k=1}^N p_k \nu(\tau_k^{-1}x) \quad (x \in X).$$

(Since τ_i are contractions hence injective, each set $\tau_k^{-1}x$ contains at most one point.) Since ν is a finite measure, there are only countably many atoms and they can be listed in a sequence that converges to zero. Therefore there is a biggest atom, i.e., one with maximal measure. Let x_0 be an atom such that $\nu(\{x_0\}) \geq \nu(\{x\})$ for all $x \in X$.

The sets $\tau_k^{-1}x_0$ have measure less than $\nu(\{x_0\})$ by maximality. From equation (6.4) we see that none of the sets $\tau_k^{-1}x_0$ can have measure strictly less than $\nu(\{x_0\})$. Thus $\nu(\tau_k^{-1}x_0) = \nu(\{x_0\})$, for $k \in \{1, \dots, N\}$. By induction, we obtain that $\nu(\tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1}x_0) = \nu(\{x_0\})$, for all $n \in \mathbb{N}$ and all $\omega_1, \dots, \omega_n \in \{1, \dots, N\}$. In particular, all sets $\tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1}x$ are non-empty. But, using the hypothesis, we obtain that there is an infinite number of points that have measure equal to $\nu(\{x_0\}) > 0$, and this contradicts the fact that ν is finite. $\square$

Lemma 6.5. *Let $(\tau_b)_{b \in B}$ be an affine IFS on $\mathbb{R}^d$ as before with $\#B \geq 2$. If $x \in \mathbb{R}^d$ then, for $n \in \mathbb{N}$ the set $\{\tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1}x \mid \omega_1, \dots, \omega_n \in B\}$ has at least n elements.*

Also the set $\{\tau_{\omega_n} \cdots \tau_{\omega_1}x \mid \omega_1, \dots, \omega_n \in B\}$ has at least n elements.

Proof. First, note that $\tau_{\omega_n}^{-1} \cdots \tau_{\omega_1}^{-1}x = \tau_{\eta_n}^{-1} \cdots \tau_{\eta_1}^{-1}x$ if and only if

$$\omega_1 + \cdots + R^{n-1}\omega_n = \eta_1 + \cdots + R^{n-1}\eta_n.$$

Then, take $b_1 \neq b_2$ in B , and for $i \in \{1, \dots, n\}$, define

$$\omega_k^{(i)} := \begin{cases} b_1, & \text{if } k = i, \\ b_2, & \text{if } k \neq i. \end{cases}$$

Then we cannot have for $i > j$,

$$\sum_{k=1}^n R^{k-1}\omega_k^{(i)} = \sum_{k=1}^n R^{k-1}\omega_k^{(j)},$$

because that implies $(R^{i-1} - R^{j-1})(b_1 - b_2) = 0$ so $R^{i-j}(b_1 - b_2) = (b_1 - b_2)$, (R^{j-1} is invertible). But this is false since R is expansive so R^{i-j} does not have the eigenvalue 1. We used that the scaling matrix R is invertible. But $b_1 - b_2$ cannot be an eigenvector of R^{i-j} since 1 is not in the spectrum of R . Thus the points $\tau_{\omega_n^{(i)}}^{-1} \cdots \tau_{\omega_1^{(i)}}^{-1}x$ are distinct, and this proves the first assertion. The same argument works for the second statement. $\square$

Corollary 6.6. *The strongly invariant measure of an affine IFS has no atoms.*

Proof. Follows directly from Theorem 6.4 and Lemma 6.5. $\square$

Next, we continue our study from [DuJo03] of infinite convolutions, and the corresponding infinite product formulas for the associated Hutchinson measures.

The Fourier transform of ν may be used for discerning properties of ν itself, for example via Wiener's test on the mean-square Cesaro sum. We will illustrate this in the next remark for the Bernoulli measures ν_λ . We show that the mean-square Cesaro sum is zero of high order. So this conclusion is stronger than ν_λ being non-atomic.

Remark 6.7. We can use a criterion of N. Wiener for the existence of atoms: see, e.g., [Kat04]. Wiener's theorem states that the following Cesaro limit exists and yields the sum absolute square of the atoms, i.e.,

$$(6.5) \quad \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |\hat{\nu}(t)|^2 dt = \sum_{x \in \{\text{atoms}\}} |\nu(\{x\})|^2.$$

In particular, if the expression on the left-hand side in (6.5) is zero, then there are no atoms. For the Bernoulli systems, we will show that this limit on the left is zero of exponential order.

Set

$$s(L) = \frac{1}{2L} \int_{-L}^L |\hat{\nu}(t)|^2 dt,$$

for $L \in \mathbb{R}_+$. An application of the formula for $\hat{\nu}$ in the proof of Lemma 4.9 now yields

$$(6.6) \quad s(\lambda^{-n}T) = \frac{1}{2T} \int_{-T}^T \prod_{k=1}^n \cos^2(2\pi\lambda^{-k}t) |\hat{\nu}(t)|^2 dt,$$

for all $T \in \mathbb{R}_+$, and all $n \in \mathbb{N}$. In particular, the sequence is monotone

$$s(\lambda^{-n}T) \leq \dots \leq s(\lambda^{-1}T) \leq s(T).$$

Therefore the limit $\lim_n s(\lambda^{-n}T)$ exists.

Note that, since $\lambda \in (0, 1)$, $\lambda^{-n}T \rightarrow \infty$ as $n \rightarrow \infty$. So fix T , and let n tend to infinity in (6.6): This limit therefore coincides with the limit (6.5). In fact, we show below that the limit in (6.6) is $\mathcal{O}(2^{-n})$.

Recall further from Example 3.2, that $\nu = \nu_\lambda$ is supported in the closed interval $[-\frac{1}{1-\lambda}, \frac{1}{1-\lambda}]$, so the transform $t \mapsto \hat{\nu}(t)$ is analytic, in particular C^∞ .

The product under the integral sign in (6.6) is

$$(6.7) \quad \prod_{k=1}^n \cos^2(2\pi\lambda^{-k}t) = 2^{-n} \prod_{k=1}^n (1 + \cos(4\pi\lambda^{-k}t)).$$

But $\prod_{k=1}^n (1 + \cos(4\pi\lambda^{-k}t))$ is a Riesz-product, which by [Kat04, Sections 1.3-1.4] represents a distribution D of compact support. In particular

$$\lim_{n \rightarrow \infty} \int_{-T}^T \prod_{k=1}^n (1 + \cos(4\pi\lambda^{-k}t)) |\hat{\nu}(t)|^2 dt$$

is finite, i.e., it is $D(|\hat{\nu}|^2)$.

Returning to (6.7), we conclude that

$$\lim_{n \rightarrow \infty} s(\lambda^{-n}T) = \mathcal{O}(2^{-n}) = 0.$$

Substituting back into Wiener's formula (6.5), the result follows.

Acknowledgements. The co-authors are pleased to acknowledge helpful discussions with Professors Keri Kornelson, David Larson, K.-S. Lau, Kathy Merrill, Paul Muhly, Roger Nussbaum, Judy Packer, Steen Pedersen, and Yang Wang.

REFERENCES

- [ALTW04] Aldroubi, Akram; Larson, David; Tang, Wai-Shing; Weber, Eric Geometric aspects of frame representations of abelian groups. *Trans. Amer. Math. Soc.* 356 (2004), no. 12, 4767–4786.
- [AnIl01] R. Anisca, M. Ilie, *A technique of studying sums of central Cantor sets*. *Canad. Math. Bull.* 44 (2001), no. 1, 12–18.
- [Ba00] V. Baladi, *Positive transfer operators and decay of correlations*, World Sci, Singapore 2000.
- [BKS91] T. Bedford, M. Keane and C. Series, eds.; *Ergodic theory, symbolic dynamics, and hyperbolic spaces. Papers from the Workshop on Hyperbolic Geometry and Ergodic Theory held in Trieste, April 17–28, 1989*. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1991. xvi+369 pp.
- [BoPa05] Bodmann, Bernhard G.; Paulsen, Vern I. *Frames, graphs and erasures*. *Linear Algebra Appl.* 404 (2005), 118–146.
- [BPS03] Binz, Ernst; Pods, Sonja; Schempp, Walter, *Heisenberg groups—a unifying structure of signal theory, holography and quantum information theory*. *J. Appl. Math. Comput.* 11 (2003), no. 1-2, 1–57.
- [Dau92] I. Daubechies, *Ten lectures on wavelets*. CBMS-NSF Regional Conf. Ser. in App. Math. 61. Soc. Industrial App. Math. (SIAM), Philadelphia, PA, 1992.
- [DuJo03] D.E. Dutkay, P.E.T. Jorgensen, *Wavelets on Fractals*; to appear in *Rev. Mat. Iberoamericana*, preprint 2003; <http://arxiv.org/abs/math.CA/0305443>
- [DuJo05] D.E. Dutkay, P.E.T. Jorgensen *Iterated function systems, Ruelle operators, and invariant projective measures*, preprint 2005, arxiv math.DS/0501077
- [Edg98] G.A. Edgar, *Integral, probability, and fractal measures*, Springer-Verlag, New York, 1998
- [Er39] P. Erdős, *On a family of symmetric Bernoulli convolutions*. *Amer. J. Math.* 61, (1939). 974–976.
- [Fal93] K.J. Falconer, *Wavelets, fractals and order-two densities. Wavelets, fractals, and Fourier transforms*, 39–46, *Inst. Math. Appl. Conf. Ser. New Ser.*, 43, Oxford Univ. Press, New York, 1993.
- [Ga62] A.M. Garsia, *Arithmetic properties of Bernoulli convolutions*; *Trans. Amer Math Soc* 102 (1962) 409-432.
- [Gun00] R. Gundy, *Low-pass filters, Martingales, and multiresolution analysis*, *Appl. Comput. Harmonic Anal.* 9 (2000) 204-219.
- [Hun84] T.W. Hungerford, *Algebra*, Springer-Verlag, NY 1984, first ed. 1974
- [Hut81] J.E. Hutchinson, *Fractals and selfsimilarity*, *Indiana Univ. Math. J.*, 30 (1981), 713-747.
- [JeWi35] B. Jessen, A. Wintner, *Distribution functions and the Riemann zeta function*; *Trans. Amer. Math. Soc.* 38 (1935) 48–88.
- [JoPe96] P.E.T. Jorgensen, S. Pedersen, *Harmonic analysis of fractal measures*, *Constructive Approximation*, v 12 (1996), 1-30.
- [JoPe98] P.E.T. Jorgensen, S. Pedersen, *Dense analytic subspaces in fractal L^2 -spaces*, *J. d'Analyse Math* 75, 1998, 185-228.
- [Jor04] P.E.T. Jorgensen, *Iterated function systems, representations, and Hilbert space*, *International J. Math.* 15 (2004) 813-832.
- [Jor06] Palle E. T. Jorgensen, *Analysis and Probability: Wavelets, signals, fractals*, Springer-Verlag, Graduate Texts in Math, GTM vol 234, to appear in Feb 2006.
- [Kah71] J.-P. Kahane, *Sur la distribution de certaines series aleatoires*; *Colloque de Thorie des Nombres* (Univ. Bordeaux, Bordeaux, 1969), 119–122. *Bull. Soc. Math. France*, Mm. No. 25, Soc. Math. France Paris, 1971.
- [Kat04] Y. Katznelson, *An introduction to harmonic analysis*. Third edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2004.
- [KeWi35] R. Kershner, A. Wintner; *On symmetric Bernoulli convolutions*; *Amer. J. Math.* (1935), 57, 541-548.

- [KSS03] M. Keane, K. Simon, B. Solomyak, *The dimension of graph directed attractors with overlaps on the line with an application to a problem in fractal image recognition*. Fund. Math. 180 (2003), 279–292.
- [LMW96] K.-S. Lau, M.F. Ma, J. Wang, *On some sharp regularity estimates of L^2 -scaling functions*, SIAM J. Math. Anal. 27 (1996), 835–864.
- [Nu01] R.D. Nussbaum, *Periodic points of positive linear operators and Perron-Frobenius operators*; Int. Eq. and Op. Th. 39 (2001) 41–97.
- [PeSo96] Y. Peres; B. Solomyak, *Absolute continuity of Bernoulli convolutions, a simple proof*. Math. Res. Lett. 3 (1996), 231–239.
- [PeSo00] Y. Peres, B. Solomyak, *Existence of L^q dimensions and entropy dimension for self-conformal measures*. Indiana Univ. Math. J. 49 (2000), 1603–1621.
- [PSS00] Y. Peres; W. Schlag; B. Solomyak, *Sixty years of Bernoulli convolutions; in Fractal geometry and stochastics, II* (Greifswald/Koserow, 1998), 39–65, Progr. Probab., 46, Birkhuser, Basel, 2000.
- [Rud87] W. Rudin, *Real and complex analysis*. Third edition. McGraw-Hill Book Co., New York, 1987. xiv+416 pp.
- [So95] B. Solomyak, *On the random series $\sum \pm \lambda^n$ (an Erdos problem)*. Ann. of Math. (2) 142 (1995), 611–625.
- [SSU01] K. Simon, B. Solomyak, M. Urbanski, *Invariant measures for parabolic IFS with overlaps and random continued fractions*. Trans. Amer. Math. Soc. 353 (2001), 5145–5164.
- [Wal78] P. Walters, *Equilibrium states for β -transformations and related transformations*. Math. Z. 159 (1978), no. 1, 65–88.
- [Wal01] P. Walters, *Convergence of the Ruelle operator for a function satisfying Bowen's condition*. Trans. Amer. Math. Soc. 353 (2001), no. 1, 327–347.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, 14 MACLEAN HALL, IOWA CITY, IA 52242-1419, U.S.A.,

E-mail address, Dorin Ervin Dutkay: `ddutkay@math.rutgers.edu`

E-mail address, Palle E.T. Jorgensen: `jorgen@math.uiowa.edu`